

*METHOD FOR ELIMINATING FRAGILE CELLS
FROM STORED CELLS*

Inventors: John H. Crowe et al.

Atty Docket No. 010023-000900US




FIG. 1




FIG. 2




FIG. 3




FIG. 4

METHOD FOR ELIMINATING FRAGILE CELLS
FROM STORED CELLS

Inventors: John H. Crowe et al.
Atty Docket No.: 010023-000900US




Fig. 5

Fragility index of RBCs incubated overnight at 4 or 37°C in the presence of increasing trehalose concentrations




Figure 6

METHOD FOR ELIMINATING FRAGILE CELLS
FROM STORED CELLS

Inventors: John H. Crowe et al.
Atty Docket No. 010023-000900US




Fig 7

Trehalose uptake as a function of the
concentration of the washing buffer





Fig. 8


METHOD FOR ELIMINATING FRAGILE CELLS
FROM STORED CELLS

Inventors: John H. Crowe et al.
Atty Docket No. 010023-000900US



**Control RBCs in
300 mOsm PBS**


Fig 9



**Trehalose-loaded RBCs
resuspended in 300 mM
PBS, 30 s**

R3+R4 = 27%

Fig. 10



forward scatter

**Trehalose-loaded RBCs
resuspended in 300 mM
PBS, 5 min**

**R3+R4 = 2%
of R1 events**

Fig 11




Figure 12